

Mathematical Statistics

Anna Janicka

Lecture VII, 07.04.2022

**ESTIMATOR PROPERTIES, PART III
(ASYMPTOTIC PROPERTIES)**

Plan for Today

1. Asymptotic properties of estimators
 - *asymptotic unbiasedness*
 - consistency
 - asymptotic normality
 - asymptotic efficiency

2. Consistency, asymptotic normality and asymptotic efficiency of MLE estimators



Asymptotic properties of estimators

- Limit theorems describing estimator properties when $n \rightarrow \infty$
- In practice: information on how the estimators behave for large samples, *approximately*
- Problem: usually, there is no answer to the question what sample is large enough (for the approximation to be valid)



Consistency

Let $X_1, X_2, \dots, X_n, \dots$ be an IID sample (of independent random variables from the same distribution). Let $\hat{g}(X_1, X_2, \dots, X_n)$ be a sequence of estimators of the value $g(\theta)$.

\hat{g} is a **consistent** estimator, if for all $\theta \in \Theta$, for any $\varepsilon > 0$:

$$\lim_{n \rightarrow \infty} P_{\theta}(|\hat{g}(X_1, X_2, \dots, X_n) - g(\theta)| \leq \varepsilon) = 1$$

(i.e. \hat{g} converges to $g(\theta)$ in probability)



Strong consistency

Let $X_1, X_2, \dots, X_n, \dots$ be an IID sample (of independent random variables from the same distribution). Let $\hat{g}(X_1, X_2, \dots, X_n)$ be a sequence of estimators of the value $g(\theta)$.

\hat{g} is **strong consistent**, if for any $\theta \in \Theta$:

$$P_{\theta} \left(\lim_{n \rightarrow \infty} \hat{g}(X_1, X_2, \dots, X_n) = g(\theta) \right) = 1$$

(i.e. \hat{g} converges to $g(\theta)$ almost surely)



Consistency – note

From the Glivenko-Cantelli theorem it follows that empirical CDFs converge almost surely to the theoretical CDF. Therefore, we should expect (strong) consistency from all sensible estimators.

Consistency = minimal requirement for a sensible estimator.



Consistency – how to verify?

- From the definition: for example with the use of a version of the Chebyshev inequality:

$$P(|\hat{g}(X) - g(\theta)| \geq \varepsilon) \leq \frac{E(\hat{g}(X) - g(\theta))^2}{\varepsilon^2}$$

Given that the MSE of an estimator is

$$MSE(\theta, \hat{g}) = E_{\theta}(\hat{g}(X) - g(\theta))^2$$

we get a sufficient condition for consistency:

$$\lim_{n \rightarrow \infty} MSE(\theta, \hat{g}) = 0$$

- From the LLN



Consistency – examples

- For any family of distributions with an expected value: the sample mean \bar{X}_n is a consistent estimator of the expected value $\mu(\theta) = E_{\theta}(X_1)$. Convergence from the SLLN.
- For distributions having a variance:
$$S_n^2 = \frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2 \quad \text{and} \quad \hat{S}_n^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2$$
are consistent estimators of the variance $\sigma^2(\theta) = \text{Var}_{\theta}(X_1)$. Convergence from the SLLN.

Consistency – examples/properties

- An estimator may be unbiased but inconsistent; e.g. $T_n(X_1, X_2, \dots, X_n) = X_1$ as an estimator of $\mu(\theta) = E_\theta(X_1)$.
- An estimator may be biased but consistent; e.g. the biased estimator of the variance or any unbiased consistent estimator $+ 1/n$.



Asymptotic normality

$\hat{g}(X_1, X_2, \dots, X_n)$ is an **asymptotically normal** estimator of $g(\theta)$, if for any $\theta \in \Theta$ there exists $\sigma^2(\theta)$ such that, when $n \rightarrow \infty$

$$\sqrt{n}(\hat{g}(X_1, X_2, \dots, X_n) - g(\theta)) \xrightarrow{D} N(0, \sigma^2(\theta))$$

Convergence in distribution, i.e. for any a

$$\lim_{n \rightarrow \infty} P_{\theta} \left(\frac{\sqrt{n}}{\sigma(\theta)} (\hat{g}(X_1, X_2, \dots, X_n) - g(\theta)) \leq a \right) = \Phi(a)$$

in other words, the distribution of $\hat{g}(X_1, X_2, \dots, X_n)$ is for large n similar to $N(g(\theta), \frac{\sigma^2}{n})$



Asymptotic normality – properties

- An asymptotically normal estimator is consistent (not necessarily strongly).
- A *similar* condition to unbiasedness – the expected value of the asymptotic distribution equals $g(\theta)$ (but the estimator *does not need to be* unbiased).
- **Asymptotic variance** defined as $\sigma^2(\theta)$
or $\sigma^2(\theta)/n$ – the variance of the asymptotic distribution



Asymptotic normality – what it is not

- For an asymptotically normal estimator we usually have:

$$E_{\theta} \hat{g}(X_1, X_2, \dots, X_n) \xrightarrow{n \rightarrow \infty} g(\theta)$$

$$n \operatorname{var} \hat{g}(X_1, X_2, \dots, X_n) \xrightarrow{n \rightarrow \infty} \sigma^2(\theta)$$

but these properties needn't hold, because convergence in distribution does not imply convergence of moments.



Asymptotic normality – example

- Let $X_1, X_2, \dots, X_n, \dots$ be an IID sample from a distribution with mean μ and variance σ^2 . On the base of the CLT, for the sample mean we have

$$\sqrt{n}(\bar{X} - \mu) \xrightarrow{D} N(0, \sigma^2)$$

In this case the asymptotic variance, σ^2/n , is equal to the estimator variance.



Asymptotic normality – how to prove it

In many cases, the following is useful:

Delta Method. Let T_n be a sequence of random variables such that for $n \rightarrow \infty$ we have

$$\sqrt{n}(T_n - \mu) \xrightarrow{D} N(0, \sigma^2)$$

and let $h: \mathbb{R} \rightarrow \mathbb{R}$ be a function differentiable at point μ such that $h'(\mu) \neq 0$. Then

$$\sqrt{n}(h(T_n) - h(\mu)) \xrightarrow{D} N(0, \sigma^2 (h'(\mu))^2)$$

μ, σ^2 are functions of θ

usually used when estimators are functions of statistics T_n , which can be easily shown to converge on the base of CLT



Asymptotic normality – examples cont.

In an exponential model: $MLE(\lambda) = \frac{1}{\bar{X}}$

From CLT, we get

$$\sqrt{n}(\bar{X} - \frac{1}{\lambda}) \xrightarrow{D} N(0, \frac{1}{\lambda^2})$$

so from the Delta Method for $h(t)=1/t$:

$$\sqrt{n}(\frac{1}{\bar{X}} - \lambda) \xrightarrow{D} N(0, \frac{1}{\lambda^2} \cdot (-\frac{1}{(1/\lambda)^2})^2)$$

so $\frac{1}{\bar{X}}$ is an asymptotically normal (and consistent) estimator of λ .



Asymptotic efficiency

For an asymptotically normal estimator $\hat{g}(X_1, X_2, \dots, X_n)$ of $g(\theta)$ we define **asymptotic efficiency** as

$$\text{as.ef}(\hat{g}) = \frac{(g'(\theta))^2 n}{\sigma^2(\theta) \cdot I_n(\theta)},$$

where $\sigma^2(\theta)/n$ is the asymptotic variance, i.e. for $n \rightarrow \infty$

$$\sqrt{n}(\hat{g}(X_1, X_2, \dots, X_n) - g(\theta)) \xrightarrow{D} N(0, \sigma^2(\theta))$$

modification of the definition of efficiency
to the limit case, with the asymptotic
variance in place of the normal variance

$$\text{as.ef}(\hat{g}) = \frac{(g'(\theta))^2}{\sigma^2(\theta) \cdot I_1(\theta)}$$

Relative asymptotic efficiency

Relative asymptotic efficiency for asymptotically normal estimators

$\hat{g}_1(X)$ and $\hat{g}_2(X)$

$$\text{as.ef}(\hat{g}_1, \hat{g}_2) = \frac{\sigma_2^2(\theta)}{\sigma_1^2(\theta)} = \frac{\text{as.ef}(\hat{g}_1)}{\text{as.ef}(\hat{g}_2)}$$

Note. A less (asymptotically) efficient estimator may have other properties, which will make it preferable to a more efficient one.



Relative asymptotic efficiency – examples.

Is the mean better than the median?



Relative asymptotic efficiency – examples.

Is the mean better than the median?

Depends on the distribution!

a) normal model $N(\mu, \sigma^2)$:

σ^2 known

$$\sqrt{n}(\bar{X} - \mu) \xrightarrow{D} N(0, \sigma^2)$$

$$\text{as.ef}(\hat{m}, \bar{X}) = \frac{2}{\pi} < 1$$

$$\sqrt{n}(\hat{m} - \mu) \xrightarrow{D} N(0, \frac{\pi\sigma^2}{2})$$

b) Laplace model $\text{Lapl}(\mu, \lambda)$

λ known

$$\sqrt{n}(\bar{X} - \mu) \xrightarrow{D} N(0, \frac{2}{\lambda^2})$$

$$\text{as.ef}(\hat{m}, \bar{X}) = 2 > 1$$

$$\sqrt{n}(\hat{m} - \mu) \xrightarrow{D} N(0, \frac{1}{\lambda^2})$$

c) some distributions do not have a mean...

Theorem: For a sample from a continuous distribution with density $f(x)$, the sample median is an asymptotically normal estimator for the median m

(provided the density is continuous and $\neq 0$ at point m):

$$\sqrt{n}(\hat{m} - m) \xrightarrow{D} N(0, \frac{1}{4(f(m))^2})$$



Consistency of ML estimators

Let $X_1, X_2, \dots, X_n, \dots$ be a sample from a distribution with density $f_\theta(x)$. If $\Theta \subseteq \mathbb{R}$ is an open set, and:

- all densities f_θ have the same support;
- the equation $\frac{d}{d\theta} \ln L(\theta) = 0$ has exactly one solution, $\hat{\theta}$.

Then $\hat{\theta}$ is the $MLE(\theta)$ and it is consistent

Note. MLE estimators do not have to be unbiased!



Asymptotic normality of ML estimators

Let $X_1, X_2, \dots, X_n, \dots$ be a sample with density $f_\theta(x)$, such that $\Theta \subseteq \mathbb{R}$ is open, and $\hat{\theta}$ is a consistent m.l.e. (for example, fulfills the assumptions of the previous theorem), and

- $\frac{d^2}{d\theta^2} \ln L(\theta)$ exists
- Fisher Information may be calculated, $0 < I_1(\theta) < \infty$
- the order of integration with respect to x and derivation with respect to θ may be changed

then $\hat{\theta}$ is asymptotically normal and

$$\sqrt{n}(\hat{\theta} - \theta) \xrightarrow{D} N(0, \frac{1}{I_1(\theta)})$$



Asymptotic normality of ML estimators

Additionally, if $g: \mathbb{R} \rightarrow \mathbb{R}$ is a function differentiable at point θ , such that $g'(\theta) \neq 0$, and $\hat{g}(X_1, X_2, \dots, X_n)$ is $MLE(g(\theta))$, then

$$\sqrt{n}(\hat{g}(X_1, X_2, \dots, X_n) - g(\theta)) \xrightarrow{D} N(0, \frac{(g'(\theta))^2}{I_1(\theta)})$$



Asymptotic efficiency of ML estimators

If the assumptions of the previous theorems are fulfilled, then the ML estimator (of θ or $g(\theta)$) is asymptotically efficient.



Asymptotic normality and efficiency of ML estimators – examples

- In the normal model: the mean is an asymptotically efficient estimator of μ
- In the Laplace model: the median is an asymptotically efficient estimator of μ



Summary: basic (point) estimator properties

- ☐ bias
- ☐ variance
- ☐ MSE
- ☐ efficiency

- ☐ asymptotic unbiasedness
- ☐ consistency
- ☐ asymptotic normality
- ☐ asymptotic efficiency

