

Mathematical Statistics

Anna Janicka

Lecture XI, 12.05.2022

**HYPOTHESIS TESTING III:
LR TEST FOR COMPOSITE HYPOTHESES
EXAMPLES OF ONE-SAMPLE TESTS**

Plan for today

1. LR test for composite hypotheses
2. Examples of LR tests:
 - Model I: One- and two-sided tests for the mean in the normal model, σ^2 known
 - Model II: One- and two-sided tests for the mean in the normal model, σ^2 unknown
 - + One- and two-sided tests for the variance
 - Model III: Tests for the mean, large samples
 - Model IV: Tests for the fraction, large samples



Testing simple hypotheses – reminder

We observe X . We want to test

$$H_0: \theta = \theta_0 \text{ against } H_1: \theta = \theta_1.$$

(two simple hypotheses)

We can write it as:

$$H_0: X \sim f_0 \text{ against } H_1: X \sim f_1,$$

where f_0 and f_1 are *densities* of distributions defined by θ_0 and θ_1 (i.e. P_0 and P_1)



Likelihood ratio test for simple hypotheses. Neyman-Pearson Lemma – reminder

$H_0: X \sim f_0$ against $H_1: X \sim f_1$

Let

$$C^* = \left\{ x \in X : \frac{f_1(x)}{f_0(x)} > c \right\}$$

such that $P_0(C^*) = \alpha$ and $P_1(C^*) = 1 - \beta$

Then, for any $C \subseteq \mathcal{X}$:

if $P_0(C) \leq \alpha$, then $P_1(C) \leq 1 - \beta$.

(i.e.: the test with critical region C^* is the most powerful test
for testing H_0 against H_1)



Neyman-Pearson Lemma – Example 1 reminder

Normal model: X_1, X_2, \dots, X_n are an IID sample from $N(\mu, \sigma^2)$, σ^2 is known

The most powerful test for

$$H_0: \mu = 0 \text{ against } H_1: \mu = 1. \quad \leftarrow \mu_0 < \mu_1$$

At significance level α :

$$C^* = \left\{ (x_1, x_2, \dots, x_n) : \bar{X} > u_{1-\alpha} \sigma / \sqrt{n} \right\}$$

If we had

$$H_0: \mu = 0 \text{ against } H_1: \mu = -1, \text{ then}$$

$$C_1^* = \left\{ (x_1, x_2, \dots, x_n) : \bar{X} < u_{1-\alpha} \sigma / \sqrt{n} \right\}$$

Neyman-Pearson Lemma – Example 1 cont.

Power of the test

$$\begin{aligned} P_1(C^*) &= P\left(\bar{X} > 1.645\sigma/\sqrt{n} \mid \mu = 1\right) = \dots \\ &= 1 - \Phi\left(1.645 - \mu_1 \cdot \sqrt{n}/\sigma\right) \approx 0.91 \end{aligned}$$

If we change α , μ_1 , n – the power of the test....



Neyman-Pearson Lemma:

Generalization of example 1

The same test is UMP for $H_1: \mu > 0$ and for
 $H_0: \mu \leq 0$ against $H_1: \mu > 0$

more generally: under additional assumptions about the family of distributions, the same test is UMP for testing

$$H_0: \mu \leq \mu_0 \text{ against } H_1: \mu > \mu_0$$

Note the change of direction of the inequality in the condition when testing

$$H_0: \mu \geq \mu_0 \text{ against } H_1: \mu < \mu_0$$



Neyman-Pearson Lemma – Example 2

Exponential model: X_1, X_2, \dots, X_n are an IID sample from $\text{distr exp}(\lambda)$, $n = 10$.

MP test for

$H_0: \lambda = 1/2$ against $H_1: \lambda = 1/4$.

At significance level $\alpha = 0.05$:

$$C^* = \left\{ (x_1, x_2, \dots, x_{10}) : \sum x_i > 31.41 \right\}$$

E.g. for a sample: 2; 0.9; 1.7; 3.5; 1.9; 2.1; 3.7; 2.5; 3.4; 2.8:
 $\Sigma = 24.5 \rightarrow$ no grounds for rejecting H_0 .



Neyman-Pearson Lemma – Example 2'

Exponential model: X_1, X_2, \dots, X_n are an IID sample from $\text{distr exp}(\lambda)$, $n = 10$.

MP test for

$H_0: \lambda = 1/2$ against $H_1: \lambda = 3/4$.

At significance level $\alpha = 0.05$:

$$C^* = \left\{ (x_1, x_2, \dots, x_{10}) : \sum x_i < 10.85 \right\}$$

E.g. for a sample: 2; 0.9; 1.7; 3.5; 1.9; 2.1; 3.7; 2.5; 3.4; 2.8:
 $\Sigma = 24.5 \rightarrow$ no grounds for rejecting H_0 .



Example 2 cont.

The test

$$C^* = \left\{ (x_1, x_2, \dots, x_{10}) : \sum x_i > 31.41 \right\}$$

is UMP for $H_0: \lambda \geq 1/2$ against $H_1: \lambda < 1/2$

The test

$$C^* = \left\{ (x_1, x_2, \dots, x_{10}) : \sum x_i < 10.85 \right\}$$

is UMP for $H_0: \lambda \leq 1/2$ against $H_1: \lambda > 1/2$

Likelihood ratio test for composite hypotheses

$X \sim P_\theta$, $\{P_\theta: \theta \in \Theta\}$ – family of distributions

We are testing $H_0: \theta \in \Theta_0$ against $H_1: \theta \in \Theta_1$

such that $\Theta_0 \cap \Theta_1 = \emptyset$, $\Theta_0 \cup \Theta_1 = \Theta$

Let

$H_0: X \sim f_0(\theta_0, \cdot)$ for some $\theta_0 \in \Theta_0$.

$H_1: X \sim f_1(\theta_1, \cdot)$ for some $\theta_1 \in \Theta_1$,

where f_0 and f_1 are densities (for $\theta \in \Theta_0$ and $\theta \in \Theta_1$, respectively)

Likelihood ratio test for composite hypotheses – cont.

$$\text{Test statistic: } \lambda = \frac{\sup_{\theta_1 \in \Theta_1} f_1(\theta_1, X)}{\sup_{\theta_0 \in \Theta_0} f_0(\theta_0, X)}$$

$$\text{or } \lambda = \frac{f_1(\hat{\theta}_1, X)}{f_0(\hat{\theta}_0, X)}$$

where $\hat{\theta}_0, \hat{\theta}_1$ are MLE for the null and alternative hypothesis models

We reject H_0 if $\lambda > c$ for a constant c
(determined according to significance level)



Likelihood ratio test for composite hypotheses

– justification

Just like in the Neyman-Pearson Lemma, we compare the “highest chance of obtaining observation X , when the alternative is true” to the “highest chance of obtaining observation X , when the null is true”; we reject the null hypothesis in favor of the alternative if this ratio is very unfavorable for the null.



Likelihood ratio test for composite hypotheses – alternative version

Test statistic:
$$\tilde{\lambda} = \frac{\sup_{\theta \in \Theta} f(\theta, X)}{\sup_{\theta_0 \in \Theta_0} f_0(\theta_0, X)}$$

or
$$\tilde{\lambda} = \frac{f(\hat{\theta}, X)}{f_0(\hat{\theta}_0, X)}$$

where $\hat{\theta}, \hat{\theta}_0$ are the ML estimators for the model without restrictions and for the null model, respectively.

We reject H_0 if $\tilde{\lambda} > \tilde{c}$ for a constant \tilde{c} .



Likelihood ratio test for composite hypotheses

– properties

For some models with composite hypotheses the UMPT *does not exist* (so the LR test will not be UMP because there is no such test)

e.g. testing $H_0: \theta = \theta_0$ against $H_1: \theta \neq \theta_0$ if the family of distributions has a *monotonic LR property*, i.e. $f_1(x)/f_0(x)$ is an increasing function of a statistic $T(x)$ for any f_0 and f_1 corresponding to parameters $\theta_0 < \theta_1$.

In order to have UMPT for $H_0: \theta = \theta_0$ against $H_1: \theta > \theta_0$ we would need a critical region of the type $T(x) > c$, and to have a UMPT for $H_0: \theta = \theta_0$ against $H_1: \theta < \theta_0$ we would need a critical region of the type $T(x) < c$, so it is impossible to find a UMPT for $H_1: \theta \neq \theta_0$.



Likelihood ratio test: special cases

The exact form of the test depends on the distribution.

In many cases, finding the distribution is hard/complicated (in many such cases, we use the asymptotic properties of the LR test instead of precise formulae)



Notation

$x_{\text{something}}$ **always** means a quantile of rank something



Model I: comparing the mean

Normal model: X_1, X_2, \dots, X_n are an IID sample from $N(\mu, \sigma^2)$, where σ^2 is **known**

$$H_0: \mu = \mu_0$$

Test statistic:

$$U = \frac{\bar{X} - \mu_0}{\sigma} \sqrt{n} \sim N(0,1)$$

$$H_0: \mu = \mu_0 \text{ against } H_1: \mu > \mu_0$$

$$\text{critical region } C^* = \{x : U(x) > u_{1-\alpha}\}$$

$$H_0: \mu = \mu_0 \text{ against } H_1: \mu < \mu_0$$

$$\text{critical region } C^* = \{x : U(x) < u_\alpha = -u_{1-\alpha}\}$$

$$H_0: \mu = \mu_0 \text{ against } H_1: \mu \neq \mu_0$$

$$\text{critical region } C^* = \{x : |U(x)| > u_{1-\alpha/2}\}$$

Model I: example

Let X_1, X_2, \dots, X_{10} be an IID sample from $N(\mu, 1^2)$:

-1.21 -1.37 0.51 0.37 -0.75 0.44 1.20 -0.96 -1.14 -1.40

Is $\mu = 0$? (for $\alpha = 0.05$)

In the sample: mean = -0.43, ~~variance = 0.92~~

Test statistic:
$$U = \frac{-0.43 - 0}{1} \sqrt{10} \approx -1.36$$

$H_0: \mu = 0$ against $H_1: \mu \neq 0$, $u_{0.975} \approx 1.96$ (p-value ≈ 0.172)

$H_0: \mu = 0$ against $H_1: \mu < 0$, $u_{0.05} \approx -1.64$ (p-value ≈ 0.086)

$H_0: \mu = 0$ against $H_1: \mu > 0$, $u_{0.95} \approx 1.64$ (p-value ≈ 0.914)

→ in none of these cases are there grounds to reject H_0 for $\alpha = 0.05$

→ but we would reject $H_0: \mu = 0$ in favor of $H_1: \mu < 0$ for $\alpha = 0.1$

Model II: comparing the mean

Normal model: X_1, X_2, \dots, X_n are an IID sample from $N(\mu, \sigma^2)$, where σ^2 is **unknown**

$$H_0: \mu = \mu_0$$

$$T = \frac{\bar{X} - \mu_0}{S} \sqrt{n} \sim t(n-1)$$

Test statistic:

$$H_0: \mu = \mu_0 \text{ against } H_1: \mu > \mu_0$$

critical region

$$C^* = \{x : T(x) > t_{1-\alpha}(n-1)\}$$

$$H_0: \mu = \mu_0 \text{ against } H_1: \mu < \mu_0$$

critical region

$$C^* = \{x : T(x) < t_{\alpha}(n-1)\}$$

$$H_0: \mu = \mu_0 \text{ against } H_1: \mu \neq \mu_0$$

critical region

$$C^* = \{x : |T(x)| > t_{1-\alpha/2}(n-1)\}$$



$$t_{\alpha}(n-1) = -t_{1-\alpha}(n-1)$$

Model II: example (mean)

Let X_1, X_2, \dots, X_{10} be an IID sample from $N(\mu, \sigma^2)$:

-1.21 -1.37 0.51 0.37 -0.75 0.44 1.20 -0.96 -1.14 -1.40

Is $\mu = 0$? (for $\alpha = 0.05$)

In the sample: mean = -0.43, variance = 0.92

Test statistic:
$$U = \frac{-0.43 - 0}{\sqrt{0.92}} \sqrt{10} \approx -1.42$$

$H_0: \mu = 0$ vs $H_1: \mu \neq 0$, $t_{0.975}(9) \approx 2.26$ (p-value ≈ 0.188)

$H_0: \mu = 0$ vs $H_1: \mu < 0$, $t_{0.05}(9) \approx -1.83$ (p-value ≈ 0.094)

$H_0: \mu = 0$ vs $H_1: \mu > 0$, $t_{0.95}(9) \approx 1.83$ (p-value ≈ 0.906)

→ in none of these cases are there grounds to reject H_0 for $\alpha = 0.05$

→ but we would reject $H_0: \mu = 0$ in favor of $H_1: \mu < 0$ for $\alpha = 0.1$

Model II: comparing the variance

Normal model: X_1, X_2, \dots, X_n are an IID sample from $N(\mu, \sigma^2)$, where σ^2 is **unknown**

$$H_0: \sigma = \sigma_0$$

$$\chi^2 = \frac{(n-1)S^2}{\sigma_0^2} \sim \chi^2(n-1)$$

Test statistic:

$$H_0: \sigma = \sigma_0 \text{ against } H_1: \sigma > \sigma_0$$

critical region

$$C^* = \{x : \chi^2(x) > \chi_{1-\alpha}^2(n-1)\}$$

$$H_0: \sigma = \sigma_0 \text{ against } H_1: \sigma < \sigma_0$$

critical region

$$C^* = \{x : \chi^2(x) < \chi_{\alpha}^2(n-1)\}$$

$$H_0: \sigma = \sigma_0 \text{ against } H_1: \sigma \neq \sigma_0$$

critical region

$$C^* = \{x : \chi^2(x) < \chi_{\alpha/2}^2(n-1) \vee \chi^2(x) > \chi_{1-\alpha/2}^2(n-1)\}$$



Model II: example (variance)

Let X_1, X_2, \dots, X_{10} be an IID sample from $N(\mu, \sigma^2)$:

-1.21 -1.37 0.51 0.37 -0.75 0.44 1.20 -0.96 -1.14 -1.40

Is $\sigma=1$? (for $\alpha = 0.05$)

In the sample: variance = 0.92

Test statistic:

$$\chi^2 = \frac{9 \cdot 0.92}{1} \approx 8.28$$

$H_0: \sigma = 1$ against $H_1: \sigma > 1$ $\chi_{0.95}^2 \approx 16.92$

$H_0: \sigma = 1$ against $H_1: \sigma < 1$ $\chi_{0.05}^2 \approx 3.33$

$H_0: \sigma = 1$ against $H_1: \sigma \neq 1$ $\chi_{0.025}^2 \approx 2.70$; $\chi_{0.975}^2 \approx 19.02$

→ in none of these cases are there grounds to reject H_0 (for $\alpha = 0.05$)



Model III: comparing the mean

Asymptotic model: X_1, X_2, \dots, X_n are an IID sample from a distribution with mean μ and variance (unknown), n – large.

$$H_0: \mu = \mu_0$$

Test statistic:

$$T = \frac{\bar{X} - \mu_0}{S} \sqrt{n}$$

has, for large n , an *approximate* distribution $N(0,1)$

$$H_0: \mu = \mu_0 \text{ against } H_1: \mu > \mu_0$$

critical region

$$C^* = \{x : T(x) > u_{1-\alpha}\}$$

$$H_0: \mu = \mu_0 \text{ against } H_1: \mu < \mu_0$$

critical region

$$C^* = \{x : T(x) < u_\alpha = -u_{1-\alpha}\}$$

$$H_0: \mu = \mu_0 \text{ against } H_1: \mu \neq \mu_0$$

critical region

$$C^* = \{x : |T(x)| > u_{1-\alpha/2}\}$$



Model IV: comparing the fraction

Asymptotic model: X_1, X_2, \dots, X_n are an IID sample from a two-point distribution, n – large.

$$P_p(X = 1) = p = 1 - P_p(X = 0)$$

$H_0: p = p_0$

Test statistic:
$$U^* = \frac{\bar{X} - p_0}{\sqrt{p_0(1 - p_0)}} \sqrt{n} = \frac{\hat{p} - p_0}{\sqrt{p_0(1 - p_0)}} \sqrt{n}$$

has an *approximate* distribution $N(0,1)$ for large n

$H_0: p = p_0$ against $H_1: p > p_0$
critical region $C^* = \{x : U^*(x) > u_{1-\alpha}\}$

$H_0: p = p_0$ against $H_1: p < p_0$
critical region $C^* = \{x : U^*(x) < u_\alpha = -u_{1-\alpha}\}$

$H_0: p = p_0$ against $H_1: p \neq p_0$
critical region $C^* = \{x : |U^*(x)| > u_{1-\alpha/2}\}$



Model IV: example

We toss a coin 400 times. We get 180 heads. Is the coin symmetric?

$$H_0: p = 1/2 \quad U^* = \frac{(180/400 - 1/2)}{\sqrt{1/2(1 - 1/2)}} \sqrt{400} = -2$$

for $\alpha = 0.05$ and $H_1: p \neq 1/2$ we have $u_{0.975} = 1.96 \rightarrow$ we reject H_0

for $\alpha = 0.05$ and $H_1: p < 1/2$ we have $u_{0.05} = -u_{0.95} = -1.64$

\rightarrow we reject H_0

for $\alpha = 0.01$ and $H_1: p \neq 1/2$ we have $u_{0.995} = 2.58$

\rightarrow we do not reject H_0

for $\alpha = 0.01$ and $H_1: p < 1/2$ we have $u_{0.01} = -u_{0.99} = -2.33$

\rightarrow we do not reject H_0

p-value for $H_1: p \neq 1/2$: 0.044

p-value for $H_1: p < 1/2$: 0.022



